

Evaluation of Hematological Changes in The Pregnant Women in EL-Beyda City- Libya

Najat M. H. Mohammed¹, Kasem A. Farag² and Hanan H. Ahmed³

¹Faculty of Medical Technology, Laboratory Medicine Department, Omer AL-Mukhtar University, EL-Beyda-Libya

²Faculty of Science, Mathematics Department, Omer AL-Mukhtar University, EL-Beyda-Libya

³Gynecology Department , Diabetic Center AL-Jabal AL-Akhdar, EL-Beyda Libya

Article information	Abstract
<p>Key words Pregnant women, Anemia, Trimester, Parity and Gravida.</p> <p>Received 2024/01 /03, Accepted 2024/01 /10, Available online /01 2024/01</p>	<p>Anemia in pregnancy is considered a major public health problem and is commonly associated with many consequences as low birth weight, premature delivery and the increased risk of the maternal mortality and morbidity in the pregnant women, mainly in the developing countries. The present study was designed to estimate the changes in hematological parameters in women during pregnancy at EL-Bayda city. Cross sectional study involved 350 randomly selected pregnant mothers who are attended several medical clinics in the period of February to July 2022, of age group (17-48) years. A venous blood samples were collected in EDTA tubes and analyzed for complete blood counts (CBC) to evaluate hemoglobin levels (Hb), mean cell hemoglobin (MCV), mean cell hemoglobin (MCH), mean cell hemoglobin concentration (MCHC), white blood cells (WBCs) and platelets count. Other data were also reported including women age, trimester of pregnancy, the number of the gravida and the parity. Results will be analyzed by SPSS version 26 and represented as mean \pm SE and P value is considered significant when it is less than 0.05. The mean age of the participants women was 29.5(\pm0.33), the overall prevalence of anemia was 57% and vast of them were in third trimester period. Out of 198 anemic women, 91 (45%) of them were suffered from microcytic hypochromic anemia (mean of MCV 71.5fl) while the rest (97) had normocytic normochromic anemia (Mean of MCV 88fl). The present investigation was revealed that the mean of hemoglobin of anemic women was (9.83\pm0.062) which are significantly difference in the mean of non anemic women (p value =0.000) as well as majority of women 165 (81 %) had mild anemia (Hb= 9-10.9 g/dl). We can conclude that the prevalence of anemia in Libya is considerably high in comparison to other countries, therefore the great awareness efforts are recommended to encourage a regular antenatal checkups.</p>

I- INTRODUCTION

Anemia in pregnancy is considered a major public health problem and still has a negative impact contributing to the maternal, neonatal mortality and morbidity around the world [1-3], affecting 35-75% of pregnant women in Africa [4, 5]. According to WHO anemia in pregnant women is diagnosed when hemoglobin concentration (less than 11g/dl) or hematocrit value (HCT) are less than 33% [6] The reduction in hemoglobin (Hb) contents inside the red blood cells will diminish oxygen-carrying capacity to the maternal body tissues [7]. Hematological parameters are usually changed physiologically during pregnancy which are related to the increase of the body demand to the nutritional factors such as folic acid,

vitamin B12 and iron as fetus requirements increase as well [8, 9]. One of these changes is the increase of total blood volume by 1.5 Liters to nourish the new fetus vascular bed demand [7, 10, 11]. In addition there is a drop in hemoglobin content about 1-2 g/dl by late of the second trimester and then stabilize in the third trimester [11, 12]. The recent investigations have reported that almost 600,000 women between the ages of 15 and 44 years worldwide are known to die each years as a result of pregnancy's and childbirth complications [13]. The proportion of maternal deaths due to anemia as estimated in some countries such as India (16%), Kenya (11%), Nigeria (9%) and Malawi (8%) [13]. In comparison with Arabic countries the prevalence was higher about third of pregnant women were suffered from anemia, 30% in Egypt, Saudi

Arabia 40% and Iraq 31% [14, 15], also the pregnant women usually suffer from leukocytosis contributed to physiological stress [11, 16]. The Previous studies have also showed that the platelet count normally decrease during pregnancy, particularly in the third trimester, this is termed as "gestational thrombocytopenia".[16, 17] The most widesprea

nutritional anemia in the pregnant women are iron deficiency anemia affecting about 80% of women population followed by folate deficiency (megaloblastic anemia) with 76% as a result of inadequate diets or are not receiving prenatal iron and folate supplements [15, 18, 19]. The present study aimed to estimate the hematological changes occurred during pregnancy in women at EL-Bayda City

II- MATERIALSAND METHODS

This descriptive cross sectional study was included 350 randomly selected pregnant women who are seeking for antenatal care at different centers in EL-Beyda city during the period from February to March 2022. A venous blood samples were collected in EDTA tubes and analyzed for complete blood counts (CBC) to evaluate hemoglobin level

(Hb) , white blood cells (WBCs), mean cell hemoglobin (MCV), Mean cell hemoglobin (MCH), mean cell hemoglobin concentration (MCHC) and platelets count. Anemia is defined by using the WHO criteria of hemoglobin values of less than 11g/dL and classified as follow [8, 20].

Mild anemia 9.0-10.9g/dl ,

Moderate anemia 7.0-9 g/dL

Severe anemia <7.0 g/dL.

MCV is used to evaluate cell volume and considered microcytic if is less than 79fl and macrocytic if more than 100fl as well as hypochromia is commented depending MCH value (less than 32pg). The data were collected by interviewing with pregnant women including age, trimester of pregnancy, the gravida and the parity. Results were analyzed by SPSS version 26 and represented as mean \pm SE and P value will be significant if it less than 0.05

III-RESULTS

The present study involved a total of 350 pregnant women who are attended the several medical clinics in AL-Bayda city. The overall prevalence rate of anemia was 57%, the mean hemoglobin of anemic women was 9.84 ± 0.06 in comparison to hemoglobin of the normal women (12.31 ± 0.07) (Table 1). The age group of the pregnant women ranged from (17 to 48) years and the majority of anemia were found in two the age group between 15-25 and 25-34 (table 2). Out of 204 anemic women, 94 of them were suffered from microcytic hypochromic anemia (Mean MCV 72fl), while the reminder appeared

with normocytic normochromic anemia (Mean MCV 88fl) table 3. Concerning to the obstetric history, Table 4 shows that there is relationship between gestational age and anemia since anemia is more prevalent in third trimester than in the first and second trimester (0.02). While regarding to parity and gravida, the result showed that there is no correlation between number of the parity and anemia ($p=0.42$) as found in table 5.

In general, there was 136 (38.8 %) of total 350 pregnant women had elevated leukocytes count (mean 14.29 ± 0.34), also 39 (11%) of total cases had thrombocytopenia (Fig. 1).

Table 1: Hematological parameters of anemic and non -anemic pregnant women in the term of Mean \pm SE.

Parameters	Mean \pm S.E		p-value
	Anemic	Non – Anemic	P< 0.05
Hb (g/dl)	9.84 ± 0.06	12.31 ± 0.07	0.00
HCT (%)	31.36 ± 0.21	39.04 ± 0.09	0.00
MCV (fl)	72.1 ± 0.56	88.86 ± 0.35	0.00
MCH (Pg)	23.58 ± 0.16	29.34 ± 0.26	0.00
MCHC g/dl	30.19 ± 0.25	33.80 ± 0.16	0.00

Table 2. Prevalence of anemia according the age group

Age	Frequency	Percentage %
15 – 24	54	27.3
25 – 34	107	54
35 – 44	36	18.2
Above 45	1	0.5

Table 3. Types of anemia found in the study according to the severity and mean cell volume (MCV), presented as percentage

Anemia	Frequency	Percentage %
Mild	165	81
Moderate	37	18
Sever	2	1
Total	198	
Microcytic anemia	94	46%
Normocytic anemia	109	53.4%
Macrocytic anemia	1	0.5%

Table 4 : The prevalence of anemia with respect to gestational period of the study population

Trimester	Anemic cases (no.)	Non anemic cases (no.)	Prevalence of anemia%	Pearson chi square (p)
First	31	15	15%	0.02
Second	43	13	21%	
Third	130	118	63.7%	
Total	204	146		

Table 5: Association between anemia, parity and gravida by chi square test

Pregnant women	Anemic	Non anemic	Total	Pearson chi square (p)
Gravida				0.42
1-3	120	74	194	
4-6	67	61	128	
7-10	17	11	28	
Parity				0.23
0-2	131	84	215	
3-5	67	56	123	
6-8	6	6	12	





Figure 1: percentage of white blood cells (left) and platelets count(right) of study population

IV- DISCUSSION

The present study demonstrated that the overall prevalence of anemia was 57% with mean hemoglobin concentration 9.84 ± 0.11 g/dl. This prevalence is considerably higher than the estimated rate in 2011 according to WHO which found that 28% of Libyan women were anemic [1]. Our prevalence is similar to other findings conducted in Nigeria and Kenya with 54.5%, 57% respectively [2]. However the estimated prevalence observed in other countries was relatively low as seen in Algeria 40.08 %, in Bahrain 26.2%, and in Macca city 39% [3]. These variations in the prevalence may be due to different methodology and sampling techniques used, nutritional habits and characteristics of the target population. In addition, 46% of anemic cases had microcytic hypochromic anemia and this may be due to iron deficiency and depletion of body iron stores which further related to the frequency of the pregnancies [2, 4]. This in agreement with the previous findings showed that about of 45-89% of pregnant

women around the world were iron deficient. A recent study in 2021 was carried in Ethiopia reported that the incidence of iron deficiency anemia was 19.3% [5]. The remaining of anemic women about 53.4% showed normocytic anemia which may refer to the presence of other causes such as urinary tract infection or blood loss. Some literatures have documented that consuming of black tea immediately after meals will reduce iron absorption due to its inhibitory effect on iron bioavailability [6], the presence of phenols in tea and coffee are considered as the potent inhibitors of iron absorption [7]. beside to the unhealthy habits of low intake of red meat, iron rich foods and vegetables was also associated with anemia (8, 9). Thus the manner of the diet during pregnancy should be investigated since the most of Libyan population drink tea immediately after meals, and it is supposed to introduce other important biochemical tests including measurement of serum iron, ferritin should to classify the types of anemia.

V-CONCLUSION

Another notable result is that the mean of hemoglobin of anemic women was 9.84 ± 0.06 g/which differ significantly to the mean of non-anemic mothers ($p=0.00$) this the drop of hemoglobin content could have a serious negative consequences on women health since the most other studies showed that anemia leads to intrauterine growth retardation, preterm delivery and low birth weight [10]. The predominant anemia in this study was in the mild stage according to WHO classification (Hb 9.9-10.9g/dl) which is different from another finding in Iraq in 2020 that found that the most of participants were in the moderate stage and some in the sever level.

Our finding showed that most of anemic cases where in third trimester period indicating the increased demand for fetal growth and accelerated of red cell production which in agreement with some publications who said that the prevalence of anemia is higher in third and second trimester than in the first trimester [3, 11]. Similarly, another study was conducted in Ethiopia found that prevalence were common in second and third trimester [3, 10]. Our study showed that there was no association between anemia and number of pregnancies ($p=0.42$), which is contrary to other reports who observed that pregnant women with multiple parity are more likely to have anemia than those with one parity, which is contributed to the reduced of maternal iron reserves in prior pregnancies and lactations periods [3, 11, 12]. One of the interesting findings is that of total 350 pregnant women, 136 (39%) of them appeared with leukocytosis mainly neutrophilia indicating of infection. However some reporters explained this elevation as physiological condition related to impaired neutrophil apoptosis as a result of pregnancy stress [13,14]. Moreover, others reported that monocytes counts usually elevated during pregnancy especially in the first trimester due to its role in preventing fetal allograft rejection by infiltrating the decidual tissue (7th–20th week of gestation) [13].

Early visit to antenatal care center during first trimester of pregnancies could strongly effective to prevent anemia intervention through regularly iron supplementations [11]

We can conclude that anemia in pregnant women is associated with negative maternal outcomes therefore, great efforts should be put in order to prevent and manage the anemia during pregnancy starting from government policy to promote health education and provide supplement for mothers in antenatal care for free.

ACKNOWLEDGMENTS

The authors would like to thank Diabetes Medical Center-EL-jabal AL-Akhdar –AL-Beyda for their permission to access the Gynecology unit to collect the data.

REFERENCES

- [1] Kadhim Jasim S, Al-Momen H, Al-Asadi F. Maternal Anemia Prevalence and Subsequent Neonatal Complications in Iraq. *Open Access Macedonian Journal of Medical Sciences*. 2020;8(B):71-5.
- [2] Goodarzi E, Beiranvand R, Naemi H, Darvishi I, Khazaei Z. Prevalence of iron deficiency anemia in Asian female population and human development index (HDI): an ecological study. *Korean Society of Obstetrics and Gynecology*. 2020;63(4):497-505.
- [3] Okube OT, Mirie W, Odhiambo E, Sabina W, Habtu M. Prevalence and factors associated with anaemia among pregnant women attending antenatal clinic in the second and third trimesters at Pumwani Maternity Hospital, Kenya. *OJOG*. 2016; 6:16-27.
- [4] Hussain AM, Kadhem QI, Hussain NA. Prevalence of anaemia in a sample of pregnant women in Babylon Governorate, Iraq. *Revista Latinoamericana de Hipertension*. 2020;15(4):275-9.
- [5] Anjum A, Manzoor M, Manzoor N, Shakir HA. Prevalence of anemia during pregnancy indistrict Faisalabad Pakistan Punjab Univ J Zool. 2015;30(1):15-20.
- [6] Dessalegn FN, Wanamo TE, Wordofa D. Prevalence of Iron Deficiency Anemia and Associated Factors among Pregnant Women Attending Antenatal Care Follow up at Dodola General Hospital West Arsi Zone Oromia Region South East Ethiopia. *Archives of Medicine*. 2021;13(8):0-.
- [7] Lebso M, Anato A, Loha E. Prevalence of anemia and associated factors among pregnant women in Southern Ethiopia: A community based cross-sectional study. *PloS one*. 2017;12(12):1-11.

- [8] Al-Wousabi F, Al-Nood H, Al-Ra'awy S. Prevalence of anemia in pregnant Yemeni women in sana'a city. Sana'a University Journal of Medical Sciences. 2012;6 (1):1-5.
- [9] Berhe B, Mardu F, Legese H, Gebrewahd A, Gebremariam G, Tesfay K, et al. Prevalence of anemia and associated factors. 2019.
- [10] Young MF, Oaks BM, Tandon S, Martorell R, Dewey KG, Wendt AS. Maternal hemoglobin concentrations across pregnancy and maternal and child health: a systematic review and meta - analysis. Annals of the New York Academy of Sciences. 2019;1450(1):47-68.
- [11] Chandra S, Tripathi AK, Mishra S, Amzarul M, Vaish AK. Physiological Changes in Hematological Parameters. 2012.
- [12] Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Vitamin and Mineral Nutrition Information System Geneva: World health organization. 2011.
- [13] Hoque M, Kader SB, Hoque E. Prevalence of Anaemia in Pregnancy at Uthungulu Health district of KwaZulu-Natal, South Africa. South African Family Practice. 2007;49(6):16-d.
- [14] Elzahaf RA, Omar M. Prevalence of anaemia among pregnant women in Derna city, Libya. Int J Community Med Public Health. 2016;3(7):1915-20.
- [15] Imran ms, helmy ya, yahya m, hafez a. Anemia with pregnancy. SOHAG MEDICAL JOURNAL. 2019;23(3):100-4.
- [16] Fleming AF. Hematological changes in pregnancy. Clin Obstet Gynecol 1975;2:269.
- [17] Azab AE, Albasha MO, Jbireal JM, Hemady SYE. Haematological Changes during Pregnancy insit into anemia, leukocytosis and thrombocytopenia. East African Scholars J Med Sci. 2020;3 (5):185-92.
- [18] Fkadu Nugusu Dessalegn, Wanamo TE, Wordofa D. prevalence-of-iron-deficiency-anemia-and-associated-factors-among-pregnant-women-attending-antenatal-care-follow-up-at-dodola-gene. Archives of Medicine. 2021;13.
- [19] Ahamed NH, Kotb SA, Hassanen RH. Knowledge and attitude of pregnant women about iron deficiency anemia in Assist University Women Health Hospital, Egypt. J Nurs Health Sci. 2018;7(3):49-58.
- [20] Eweis M, Farid EZ, El-Malky N, Abdel-Rasheed M, Salem S, Shawky S. Prevalence and determinants of anemia during the third trimester of pregnancy. Clinical Nutrition ESPEN. 2021;44:194-9.
- [21] Abedi GAA, Arar AA, Alridh MSA. Assessment of Pregnant Women Knowledge and Practices Concerning Iron Deficiency Anemia at Al-Amara City/Iraq. Medico-legal Updat. 2020;20(3):152-5.
- [22] Fuzi SFA, Dagmar Koller, Sylvaine Bruggraber, Dora IA Pereira, Jack R Dainty, Mushtaq S. A 1-h time interval between a meal containing iron and consumption of tea attenuates the inhibitory effects on iron absorption: a controlled trial in a cohort of healthy UK women using a stable iron isotope. Am J Clin Nutr 2017;106:1413-21.
- [23] 23. Wolde T. Effects of caffeine on health and nutrition: A Review. Food Science and Quality Management. 2014;30:59-65.
- [24] Soliman GZA, Azmi MN, El-S S. Prevalence of Anemia in Egypt (Al-Gharbia Governorate). The Egyptian Journal of Hospital Medicine 2007;28:295-305.
- [25] Taner üE, Ekin A, Solmaz U, Erpala MB, Gezer C, Özeren M, et al. Prevalence and risk factors of anemia among pregnant women attending a high-volume tertiary care. J Turk Ger Gynecol Assoc 2015;16:231-6.
- [26] Al-Aini S, Senan CP, Azzani M. Prevalence and associated factors of anemia among pregnant women in Sana'a, Yemen. Indian Journal of Medical Sciences. 2020;72:185-90.